TEST #3 STA 5326 December 1, 2010

Name:

Please read the following directions. DO NOT TURN THE PAGE UNTIL INSTRUCTED TO DO SO

Directions

- This exam is **closed book** and **closed notes**. (You will have access to a copy of the "Table of Common Distributions" given in the back of the text.)
- The different problems are not related.
- The different parts of a problem are sometimes unrelated. If you cannot solve part of a problem, you should still go on to look at the later parts.
- Show and explain your work (including your calculations) for all problems unless you are explicitly told otherwise. No credit is given without work. But don't get carried away! Show enough work so that what you have done is clearly understandable.
- Partial credit is available. (If you know part of a solution, write it down. If you know an approach to a problem, but cannot carry it out write down this approach. If you know a useful result, write it down.)
- All the work on the exam should be your own. No "cooperation" is allowed.
- Arithmetic does **not** have to be done completely. Answers can be left as fractions or products. You do not have to evaluate binomial coefficients, factorials or large powers. Answers can be left as summations (unless there is a simple closed form such as when summing a geometric or exponential series).
- You need only pens, pencils, erasers and a calculator. (You will be supplied with scratch paper.)
- Do **not** quote homework results. If you wish to use a result from homework in a solution, you must prove this result.
- The exam has 9 pages.
- There are a total of 100 points.

The Bivariate Normal Density:

$$f(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(-\frac{1}{2(1-\rho^2)}\left(\left(\frac{x-\mu_X}{\sigma_x}\right)^2 - 2\rho\left(\frac{x-\mu_X}{\sigma_x}\right)\left(\frac{y-\mu_Y}{\sigma_Y}\right) + \left(\frac{y-\mu_Y}{\sigma_Y}\right)^2\right)\right)$$

Problem 1. Suppose X and Y have the joint density

$$f(x,y) = \frac{1}{8\pi\sqrt{1-\rho^2}} \exp\left(-\frac{1}{8(1-\rho^2)} \left(x^2 - 2\rho xy + y^2\right)\right) \quad \text{for } -\infty < x < \infty, -\infty < y < \infty$$

(a) (3%) What is the marginal distribution of X? State the name of the distribution and the values of any parameters. (No work is required for this part.)

the marginal distribution of \boldsymbol{X} is ______

(b) (9%) Verify your answer by calculating the marginal density of X from the joint density given above.

 $f_X(x) =$ _____

[Problem 1 continued]

(c) (9%) Calculate the conditional density of Y given X = x.

 $f_{Y|X}(y|x) =$ _____

(d) (3%) What is the conditional distribution of Y given X = x? Name the distribution whose density you found above, and give the values of any parameters. (No work is required.)

cond. distn. of Y given X = x is _____

Problem 2. (20%) Let (X, Y) be a bivariate random vector with joint pdf f(x, y). Let

$$S = X + aY + b, \quad T = cY + k$$

where a, b, c, k are fixed constants. Find an expression for the joint pdf of (S, T).

Problem 3. Let Y be the number of seeds in a randomly chosen packet. Suppose Y has a Poisson distribution with mean λ . Each seed, when planted, has probability p of growing and producing a plant. Suppose a packet is chosen at random and all the seeds in it are planted. Let X be the number of plants which then grow.

(a) (9%) Find the mean of X.

EX =_____

(b) (10%) Find the variance of X.

[Problem 3 continued]

(c) (4%) Are X and Y independent? Clearly answer "Yes" or "No" and justify your answer.

Problem 4. (6%) A random point (X, Y) is distributed uniformly on the **triangle** with vertices (0, 0), (1, 0), and (0, 1). For this random point, what is P(X > a) where 0 < a < 1?

Problem 5. Suppose X and Y are discrete random variables taking the values 1, 2, 3, 4. Use the table below in the following two parts.

	(Y=k)
1 1 1/20	10/20
2 4/3 3/20	6/20
3 5/3 6/20	3/20
4 2 10/20	1/20

(a) (3%) Find $P\{E(Y|X) > 1.5\}$.

 $P\left\{ E(Y|X) > 1.5 \right\} =$ ______

(b) (3%) Find $E[\{E(Y|X)\}^2]$.

Problem 6. The transition probability matrix P for a Markov chain X_0, X_1, X_2, \ldots with state space $\{1, 2, 3, 4\}$ is given below along with the matrix products P^2 and P^3 .

$$P = \begin{pmatrix} 0.11 & 0.36 & 0.23 & 0.3 \\ 0.01 & 0.03 & 0.55 & 0.41 \\ 0.07 & 0.75 & 0.14 & 0.04 \\ 0.28 & 0.19 & 0.48 & 0.05 \end{pmatrix}$$
$$P^{2} = \begin{pmatrix} 0.1158 & 0.2799 & 0.3995 & 0.2048 \\ 0.1547 & 0.4949 & 0.2926 & 0.0578 \\ 0.0362 & 0.1603 & 0.4674 & 0.3361 \\ 0.0803 & 0.476 & 0.2601 & 0.1836 \end{pmatrix}$$
$$P^{3} = \begin{pmatrix} 0.1008 & 0.3886 & 0.3348 & 0.1757 \\ 0.0586 & 0.301 & 0.3765 & 0.2639 \\ 0.1324 & 0.4322 & 0.3233 & 0.1121 \\ 0.0832 & 0.2731 & 0.4048 & 0.2388 \end{pmatrix}$$

(a) (5%) Find the value of $P_4(X_1 = 3, X_4 = 2, X_6 = 1)$. [Note: This probability may also be written as $P(X_1 = 3, X_4 = 2, X_6 = 1 | X_0 = 4)$.] (Show a little work.)

probability=

(b) (3%) Suppose the chain starts (at time zero) in a randomly chosen state with probabilities given by the vector a = (.5, .2, .2, .1). What is $P(X_{10} = 3)$, the probability the chain is in state 3 at time 10? [Use matrix notation and state your answer as a specific entry in a specific vector or matrix for which you give a formula. Do NOT attempt to evaluate your answer.] (No work is required.)

$$P(X_{10}=3) = _{-}$$

No work is required for the problems on this page. You will receive full credit just for stating (or circling) the correct answer.

Problem 7.

(a) (3%) Suppose X and Y are independent random variables with densities $f_X(x)$ and $f_Y(y)$. State a general formula for the density of Z = X + Y. (This formula is called the convolution integral.)

 $f_Z(z) =$ _____

Problem 8. (3%) Apply this formula to the case where X and Y are exponential random variables with densities $f_X(x) = \alpha e^{-\alpha x}$ for x > 0 and $f_Y(y) = \lambda e^{-\lambda y}$ for y > 0. Write an explicit integral (with explicit limits!) for the density $f_Z(z)$ of Z = X + Y. (Do NOT evaluate this integral.)

 $f_Z(z) =$ _____

Problem 9. (4%) Suppose X is a random variable with mean μ and variance σ^2 , and b is a constant. What is the value of Cov(X + b, X - b)?

a) $\sigma^2 - b^2$ b) $\sigma^2 + b^2$ c) μ^2 d) σ^2 e) $\rho\sigma^2$ f) $(\mu + b)(\mu - b)$ g) $\sigma^2 - (\mu + b)(\mu - b)$ h) 0 i) 1/2 j) 1

Problem 10. (3%) Suppose X and Z are independent random variables with $X \sim \text{Uniform}(-2, 2)$ and

$$Z = \begin{cases} +1 & \text{with probability } 1/2, \\ -1 & \text{with probability } 1/2. \end{cases}$$

Define Y = ZX. What is the value of P(|X| < 1, |Y| > 1)?

a) 0 b)
$$1/16$$
 c) $1/4$ d) $1/2$ e) $3/4$ f) 1