

Please read the following directions.

**DO NOT TURN THE PAGE UNTIL INSTRUCTED TO DO SO**

## Directions

- This exam is **closed book** and **closed notes**.
- The problems are all multiple choice or fill in the blank.
- Circle a **single** answer for each multiple choice question. Your choice should be made clearly.
- There is no penalty for guessing.
- In one problem, you need to know a fact about the normal distribution:  
 $z_{\alpha/2} = 1.96$  for  $\alpha = .05$ .
- The exam has **12** pages.
- Each multiple choice question and “fill in the blank” is worth equal credit. There are a total of 32 equally weighted items (since Problem 13 has two blanks to fill in).

**Problem 1.** You wish to apply the methods of ARIMA modeling to a time series. Suppose you observe that the variability of the series increases systematically with the level. What should you do?

- a) Try a transformation
- b) Try differencing at lag 1
- c) Try differencing at the seasonal lag
- d) Use a seasonal model
- e) Try fitting a trend
- f) None of the above

**Problem 2.** A time series has a stationary mean, but you observe that the variability increases steadily with time. Which of the following is likely to help?

- a) Try a transformation
- b) Try differencing at lag 1
- c) Try differencing at the seasonal lag
- d) Use a seasonal model
- e) Try fitting a trend
- f) None of the above

**Problem 3.** Suppose  $\{z_t\}$  is an AR(2) process:

$$z_t = C + \phi_1 z_{t-1} + \phi_2 z_{t-2} + a_t$$

If you observe a realization  $z_1, z_2, \dots, z_n$  where  $n$  is very large (say, a million), and fit a regression model

$$z_t = \beta_0 + \beta_1 z_{t-1} + \beta_2 z_{t-2} + \beta_3 z_{t-3} + \varepsilon_t$$

to this data, then the estimate of  $\beta_3$  will be approximately equal to ...

- |              |          |              |             |               |
|--------------|----------|--------------|-------------|---------------|
| a) $a_{t-3}$ | b) $C$   | c) $\phi_1$  | d) $\phi_2$ | e) $\sigma_a$ |
| f) $z_3$     | g) $a_3$ | h) $z_{t-3}$ | i) 0        | j) 1          |

**Problem 4.** Suppose  $\{z_t\}$  is a stationary AR(2) process:

$$z_t = C + \phi_1 z_{t-1} + \phi_2 z_{t-2} + a_t$$

Which one of the following pairs of random variables are independent of each other?

- |                        |                        |                        |                        |
|------------------------|------------------------|------------------------|------------------------|
| a) $z_t$ and $z_{t+1}$ | b) $z_t$ and $a_t$     | c) $z_t$ and $a_{t-1}$ | d) $z_t$ and $z_{t-2}$ |
| e) $z_t$ and $a_{t-3}$ | f) $z_t$ and $z_{t-1}$ | g) $z_t$ and $a_{t+1}$ | h) $z_t$ and $z_{t-3}$ |

**Problem 5.** In the blank provided, write an equation expressing a purely seasonal ARIMA(1, 0, 1)<sub>s</sub> model.

For problems 6 to 9, fill in the blanks with numerical answers.

These problems all use the fragment of output given below obtained from fitting a regression model  $Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \varepsilon$  for a response variable  $Y$  on two covariates  $X_1$  and  $X_2$ .

Root MSE	10.00	R-Square	0.88
Dependent Mean	12.00	Adj R-Sq	0.87
Coeff Var	20.00		

Parameter Estimates					
Variable	DF	Parameter Estimate	Standard Error	t-value	Pr >  t
Intercept	1	4.0	0.5	8.0	< .0001
X1	1	3.0	2.0	1.5	.1369
X2	1	1.0	2.5	0.4	.6900

**Problem 6.** Find a 95% confidence interval for  $\beta_1$ .

the confidence interval is \_\_\_\_\_

**Problem 7.** Assuming the residuals are approximately normally distributed with mean zero, give a value  $A$  such that you would expect about 68% of the residuals to be between  $-A$  and  $+A$ .

$A =$  \_\_\_\_\_

Consider a case with values:

$Y$	$X_1$	$X_2$
5.0	2.0	3.0

**Problem 8.** What is the **predicted value** (also known as fitted value) for this case?

predicted value= \_\_\_\_\_

**Problem 9.** What is the **residual** for this case?

residual= \_\_\_\_\_

**Problem 10.** For an  $\text{AR}(p)$  process,

- a) The theoretical **PACF** decays to zero, either exponentially or with a damped sine wave pattern or with both of these patterns.
- b) The theoretical **ACF** decays to zero, either exponentially or with a damped sine wave pattern or with both of these patterns.
- c) The theoretical **ACF** has a cutoff to zero after lag  $p$ .

**Problem 11.** A stationary  $\text{AR}(3)$  process can be re-written as a \_\_\_\_\_ process.

- a)  $\text{MA}(\infty)$       b)  $\text{AR}(2)$       c)  $\text{AR}(1)$       d)  $\text{MA}(1)$       e)  $\text{MA}(2)$       f)  $\text{MA}(3)$

**Problem 12.** Suppose  $EX = 3$  and  $EY = 7$ . What is the value of  $E(2X + 3Y + 4)$ ? (Give a numerical answer.)

$$E(2X + 3Y + 4) = \underline{\hspace{2cm}}$$

**Problem 13.** The ARMA process

$$z_t = \phi_1 z_{t-1} + \phi_2 z_{t-2} + a_t - \theta_1 a_{t-1}$$

can be written in backshift form as  $\phi(B)z_t = \theta(B)a_t$ . Write expressions for  $\phi(B)$  and  $\theta(B)$  in the blanks provided.

$$\phi(B) = \underline{\hspace{4cm}}$$

$$\theta(B) = \underline{\hspace{4cm}}$$

**Problem 14.** The ARMA process  $\phi(B)z_t = \theta(B)a_t$  can also be written as  $z_t = \psi(B)a_t$ . What is  $\psi(B)$ ?

- a)  $\phi(B)/\theta(B)$       b)  $(1 - B)^d$       c)  $1 - B$       d)  $\theta(B) + \phi(B)$
- e)  $\theta(B)/\phi(B)$       f)  $\theta(B) - \phi(B)$       g)  $(1 - B)\phi(B)$       h)  $(1 - B)\theta(B)$

**Problem 15.** Suppose  $\psi(B) = \sum_{k=0}^{\infty} \psi_k B^k$ . The expression  $\psi(B)a_t$  is equivalent to which of the following:

- a)  $a_t - \theta_1 a_{t-1} - \dots - \theta_q a_{t-q}$
- b)  $a_t + \theta_1 a_{t-1} + \dots + \theta_q a_{t-q}$
- c)  $\sum_{k=0}^{\infty} \psi_{t-k} a_t$
- d)  $\sum_{k=0}^{\infty} \psi_{t+k} a_t$
- e)  $\sum_{k=0}^{\infty} \psi_k a_{t-k}$
- f)  $\sum_{k=0}^{\infty} \psi_k a_{t+k}$

**Problem 16.** What is the formula for SBC (Schwarz's Bayesian Criterion)? (In the choices below,  $L$  is the likelihood,  $k$  is the number of parameters, and  $n$  is the number of residuals, the length of the series minus any observations lost due to differencing.)

- a)  $-2 \ln(L) + 2k$
- b)  $+2 \ln(L) - 2k$
- c)  $2L - 2k$
- d)  $-2L - 2k$
- e)  $-2 \ln(L) + k \ln(n)$
- f)  $+2 \ln(L) - k \ln(n)$

**Problem 17.** The process generated by

$$z_t = 20.0 + 0.5z_{t-1} + a_t - 1.5\theta_1 a_{t-1} + 0.8a_{t-2}$$

is called a \_\_\_\_\_ process.

- a) ARIMA(1,1,2)   b) ARIMA(2,1,1)   c) ARMA(1,2)   d) ARMA(2,1)   e) MA(2)   f) AR(1)

**Problem 18.** An MA(5) process is stationary

- a) if the roots of  $\theta(B) = 0$  lie strictly outside the unit circle
- b) if  $|\theta_1| < 1$
- c) if  $|\theta_1| > 1$
- d) if  $|\theta_2| < 1$  and  $\theta_2 \pm \theta_1 < 1$
- e) if  $|\theta_5| < 1$
- f) always
- g) never

**Problem 19.** Suppose you have fit four different non-seasonal models to a time series, and all models have acceptable residual diagnostics with all their parameters significantly different from zero. The SBC values for these models are as follows:

	Model	SBC
A	(0,0,1)	+109
B	(0,0,2)	+108
C	(2,0,0)	+107
D	(1,0,0)	+106
E	(1,0,1)	+105

Given only this information, which model is preferred?

- a) A                      b) B                      c) C                      d) D                      e) E                      f) not clear

**Problem 20.** On another time series, suppose you have fit four different seasonal models, and all models have acceptable residual diagnostics with all their parameters significantly different from zero. The AIC values for these models are as follows:

	Model	AIC
A	$(0, 0, 2)(1, 0, 0)_{12}$	-100
B	$(1, 0, 0)(1, 0, 0)_{12}$	-101
C	$(1, 1, 0)(0, 0, 1)_{12}$	-102
D	$(1, 0, 0)(1, 1, 0)_{12}$	-104

Given only this information, which model is preferred?

- a) A                      b) B                      c) C                      d) D                      e) not clear

**Problem 21.** The model

$$(1 - \phi_1 B - \phi_2 B^2 - \phi_3 B^3)(1 - \Phi_1 B^6)z_t = C + a_t$$

is a special case of a \_\_\_\_\_ model.

- a) AR(9)                      b) AR(18)                      c) AR(4)                      d) AR(6)                      e) AR(3)  
f) MA(9)                      g) MA(18)                      h) MA(4)                      i) MA(6)                      j) MA(3)

**Problem 22.** For a stationary process  $\{z_t\}$ , let  $\gamma_k = \text{Cov}(z_t, z_{t-k})$  for  $k = 0, \pm 1, \pm 2, \dots$ . Which of the following statements is always true?

- a)  $\mu_z = \frac{C}{1 - \gamma_1 - \gamma_2 - \dots - \gamma_p}$   
b)  $\gamma_k = \theta_1^k$   
c)  $\gamma_k = 0$  for  $k > q$   
d)  $\gamma_k = \frac{\sigma_a^2}{1 - \phi_k^2}$   
e)  $\rho_k = \frac{\gamma_k}{\gamma_0}$

**Problem 23.** For any stationary ARMA( $p, q$ ) process, the value of  $\phi_{kk}$  (the partial autocorrelation at lag  $k$ ) can be computed as a function of \_\_\_\_\_

- a)  $\phi_1, \phi_2, \dots, \phi_p$                       b)  $\theta_1, \theta_2, \dots, \theta_q$                       c)  $\rho_1, \rho_2, \dots, \rho_k$                       d)  $\rho_k$                       e)  $\theta_k$                       f)  $\phi_k$

**Problem 24.** The Ljung-Box statistic  $Q(m)$  is a test of the null hypothesis


- a)  $H_0 : \rho_m = 0$   
b)  $H_0 : \rho_1 = \rho_2 = \dots = \rho_m = 0$   
c)  $H_0 : \rho_1 = 0$   
d)  $H_0 : \rho_m = \rho_{m+1} = \rho_{m+2} = \dots = 0$

**Problem 25.** If the value of the Ljung-Box statistic  $Q(m)$  is large, the associated  $p$ -value will be \_\_\_\_\_.

- a) close to 1      b) close to 0      c) close to  $-1$       d) close to 0.5

**Problem 26.** The scatterplot below depicts a sample of size 100 from a population with correlation \_\_\_\_\_.

- a)  $-10$       b)  $-2$       c)  $-1$       d)  $-0.9$       e)  $-0.2$   
 f)  $0$       g)  $+0.2$       h)  $+0.9$       i)  $+1$       j)  $+2$       k)  $+10$



**Problem 27.** A list of stationary processes is given below. Which of these processes will be the most difficult to distinguish from a non-stationary process based on the sample ACF?

- a) MA(1) with  $\theta_1 = -0.5$   
 b) MA(1) with  $\theta_1 = 0.9$   
 c) MA(1) with  $\theta_1 = 0.5$   
 d) AR(1) with  $\phi_1 = -0.5$   
 e) AR(1) with  $\phi_1 = 0.9$   
 f) AR(1) with  $\phi_1 = 0.5$   
 g) random shocks

**Problem 28.** Suppose  $a_1, a_2, a_3, \dots$  is a random shock sequence. What is the value of

$$E[(\psi_0 a_3 + \psi_1 a_2 + \psi_2 a_1)^2] ?$$

- a)  $\psi_0^2 a_3^2 + \psi_1^2 a_2^2 + \psi_2^2 a_1^2$
- b)  $\sigma_a^2(\psi_0 \psi_1 + \psi_1 \psi_2)$
- c)  $\sigma_a^2(\psi_0^2 + \psi_1^2 + \psi_2^2)$
- d)  $\sigma_a(\psi_0 + \psi_1 + \psi_2)$
- e)  $\sigma_a(\psi_0 \psi_1 + \psi_1 \psi_2)$
- f)  $\sigma_z(\psi_0 + \psi_1 + \psi_2)$
- g)  $\sigma_z(\psi_0 \psi_1 + \psi_1 \psi_2)$

The following three pages (pages 9 to 11) give output (produced by the IDENTIFY statement in PROC ARIMA) for a series  $z_t$  and its first and second differences. Using this output, select a reasonable ARIMA( $p, d, q$ ) model for this series. Specify your answer in the next two questions.

**Problem 29.** What value of  $d$  should be used?

- a)  $d = 0$
- b)  $d = 1$
- c)  $d = 2$
- d)  $d = 3$

**Problem 30.** What values of  $p$  and  $q$  should be used?

- a)  $p = 1, q = 0$
- b)  $p = 2, q = 0$
- c)  $p = 3, q = 0$
- d)  $p = 0, q = 1$
- e)  $p = 1, q = 1$
- f)  $p = 0, q = 2$
- g)  $p = 0, q = 3$
- h)  $p = 2, q = 2$

**Problem 31.** The last page of the exam (page 12) gives the sample ACF and PACF of a monthly time series  $\{y_t\}$ . The time series plot is not given but appears to be stationary. Select a plausible ARIMA( $p, d, q$ )( $P, D, Q$ )<sub>12</sub> model for this series.


- a)  $(1, 0, 0)(0, 0, 1)_{12}$
- b)  $(1, 0, 0)(1, 0, 0)_{12}$
- c)  $(0, 0, 1)(0, 0, 1)_{12}$
- d)  $(0, 0, 1)(1, 0, 0)_{12}$
- e)  $(1, 1, 0)(1, 0, 0)_{12}$
- f)  $(0, 0, 1)(0, 1, 1)_{12}$
- g)  $(1, 1, 0)(0, 1, 1)_{12}$



*The ARIMA Procedure*

Name of Variable = z	
Mean of Working Series	1420.536
Standard Deviation	1251.21
Number of Observations	200


Autocorrelation Check for White Noise									
To Lag	Chi-Square	DF	Pr > ChiSq	Autocorrelations					
6	1120.89	6	<.0001	0.987	0.973	0.960	0.946	0.933	0.919
12	2084.34	12	<.0001	0.905	0.891	0.877	0.863	0.849	0.834
18	2891.39	18	<.0001	0.820	0.806	0.791	0.776	0.762	0.747
24	3548.84	24	<.0001	0.732	0.718	0.703	0.688	0.673	0.659



The ARIMA Procedure

Name of Variable = z	
Period(s) of Differencing	1
Mean of Working Series	19.6022
Standard Deviation	11.7548
Number of Observations	199
Observation(s) eliminated by differencing	1


Autocorrelation Check for White Noise									
To Lag	Chi-Square	DF	Pr > ChiSq	Autocorrelations					
6	1030.80	6	<.0001	0.981	0.957	0.927	0.900	0.875	0.853
12	1819.30	12	<.0001	0.832	0.814	0.796	0.780	0.763	0.747
18	2435.18	18	<.0001	0.730	0.713	0.695	0.677	0.659	0.642
24	2898.56	24	<.0001	0.624	0.607	0.591	0.575	0.562	0.551




The ARIMA Procedure

Name of Variable = z	
Period(s) of Differencing	1,1
Mean of Working Series	0.155902
Standard Deviation	1.551766
Number of Observations	198
Observation(s) eliminated by differencing	2

Autocorrelation Check for White Noise									
To Lag	Chi-Square	DF	Pr > ChiSq	Autocorrelations					
6	69.21	6	<.0001	0.379	0.431	-0.087	-0.029	-0.064	-0.041
12	70.80	12	<.0001	-0.017	-0.004	-0.010	-0.021	-0.066	-0.048
18	73.73	18	<.0001	-0.057	0.015	-0.039	0.040	-0.003	0.083
24	100.61	24	<.0001	0.025	-0.012	-0.115	-0.166	-0.215	-0.177



*The ARIMA Procedure*



*The ARIMA Procedure*

